Journal of Advances in Drug Research

ISSN: 2230-7761
www.jadr.co.in

Volume 17, Issue 05 | Received 2.11.2026 | Accepted 7.11.2026 | Online 12.11.2026

IMPACT OF ROOT CANAL IRRIGATION AND OBTURATION QUALITY ON CLINICAL AND RADIOGRAPHIC OUTCOMES IN CHRONIC APICAL PERIODONTITIS: AN EVIDENCE-BASED ANALYSIS

Nodir Nosirovich Shukurov

Asian International University, Bukhara, Uzbekistan
Department of Dentistry
ORCID: 0009-0003-5243-9058 | E-mail: shukurovnodir01@gmail.com

Abstract

Background: Chronic apical periodontitis is a biofilm-associated inflammatory condition for which nonsurgical root canal treatment aims to control intraradicular infection and facilitate periapical healing. The relative importance of irrigation and obturation quality must be interpreted within a multistep treatment pathway rather than as isolated technical determinants. Objective: To critically examine evidence relating irrigant selection and delivery, obturation length and homogeneity, and coronal sealing to clinical and radiographic outcomes. Methods: A targeted narrative synthesis was undertaken using published clinical guidelines, systematic reviews, meta-analyses, and observational outcome studies in endodontics. The evidence was organized by intervention, outcome, study design, and sources of bias; no primary patient data or new effect estimates were generated. Results: The 2023 European Society of Endodontology guideline supports meticulous asepsis, chemomechanical preparation, and considered use of sodium hypochlorite and EDTA, while finding insufficient outcome evidence to establish the superiority of adjunctive irrigant-activation methods. Earlier outcome syntheses associate void-free, appropriately extended fillings and adequate coronal restorations with more favorable results, although such associations are affected by baseline disease severity and procedural confounding. Comparative clinical studies have not established one obturation material or technique as consistently superior for periapical healing. Conclusion: Optimizing outcomes requires integrated infection control, technically adequate obturation, prompt definitive restoration, and structured follow-up. A prospective quality-assessment framework is proposed for future validation in patients with chronic apical periodontitis.

Keywords: chronic apical periodontitis; root canal irrigation; sodium hypochlorite; EDTA; obturation quality; root canal filling; periapical healing; coronal seal; endodontic outcomes.

Introduction

Chronic apical periodontitis is a persistent inflammatory lesion of the periradicular tissues generally associated with microbial infection of a necrotic root canal system. The clinical presentation may be relatively silent despite the presence of radiographic periapical

Journal of Advances in Drug Research

ISSN: 2230-7761
www.jadr.co.in

Volume 17, Issue 05 | Received 2.11.2026 | Accepted 7.11.2026 | Online 12.11.2026

changes, making diagnosis and post-treatment evaluation dependent on a combination of history, clinical examination, pulp testing, and imaging. Nonsurgical root canal treatment attempts to interrupt this disease process by accessing the entire canal system, reducing the microbial burden, and sealing the prepared space against reinfection. When the treated tooth is restorable, preservation through appropriately executed endodontic treatment is generally preferred to extraction [1,2].

Two technically central stages of this process are irrigation and obturation. Instrumentation cannot fully contact the irregularities of complex canal anatomy, while irrigants act through antimicrobial, tissue-dissolving, and smear-layer-removal properties. Conversely, obturation cannot compensate for inadequate infection control: its function is to occupy the prepared canal space and contribute to a durable barrier against recolonization. The quality of the definitive coronal restoration and the baseline condition of the periapical tissues further influence the probability and pace of healing [2,3].

The literature on endodontic outcomes is extensive but difficult to interpret because irrigation chemistry, activation, shaping, filling techniques, restorative procedures, and patient characteristics are interdependent. A technically dense radiographic filling may coexist with residual infection, while an initially persistent radiolucency can regress slowly despite an adequate procedure. The objective of this paper is therefore to evaluate irrigation and obturation quality as related elements of an evidence-based treatment strategy and to formulate measurable indicators for future prospective clinical research involving patients with chronic apical periodontitis.

Literature Review

The European Society of Endodontology (ESE) S3-level guideline, developed through a structured appraisal of comparative research and evidence grading, provides the principal contemporary framework for nonsurgical treatment of pulpal and apical disease. It emphasizes rigorous asepsis, dental dam isolation, adequate visibility, systematic canal preparation, restoration to function, and reassessment of healing. In asymptomatic apical periodontitis, sodium hypochlorite at concentrations of 1–5.25%, followed by EDTA and a subsequent sodium hypochlorite rinse, may be considered; however, evidence from clinical outcomes is insufficient to establish one irrigant concentration or sequence as universally superior [1,4].

Sodium hypochlorite is widely used because it combines antimicrobial activity with dissolution of organic remnants, whereas EDTA acts primarily on the inorganic component of the smear layer. Successful irrigation depends not only on the substance but also on delivered volume, fluid exchange, canal anatomy, needle position, solution renewal, and avoidance of extrusion into periapical tissues. Adjunctive procedures, including passive ultrasonic activation, laser-based approaches, photoactivated disinfection, and ozone, can alter surrogate measures of cleaning under experimental conditions. Nevertheless, improved debris removal in vitro is not interchangeable with demonstrated improvement in tooth retention or radiographic healing. The ESE recommendation does not support routine addition of these approaches when clinical outcome superiority remains unproven [1,4,5].

Journal of Advances in Drug Research

ISSN: 2230-7761
www.jadr.co.in

Volume 17, Issue 05 | Received 2.11.2026 | Accepted 7.11.2026 | Online 12.11.2026

Obturation is similarly characterized by a distinction between mechanical performance and patient-oriented outcomes. Techniques such as cold lateral compaction, warm vertical compaction, carrier-based methods, and matched-taper single-cone filling can provide acceptable results when applied with appropriate sealers and clinical control. A systematic review conducted for the ESE guideline found insufficient comparative evidence to favor one commonly evaluated filling technique or sealer category in terms of healing of apical periodontitis. Accordingly, technical proficiency, canal anatomy, safety, adequate canal disinfection, and an effective coronal seal remain more defensible selection criteria than claims of universal superiority for a particular commercial material [1,6].

Prognostic studies supply complementary evidence. In a systematic review of 63 studies, Ng and colleagues identified pre-operative periapical condition, radiographic filling length, visible voids, and the quality of coronal restoration as important factors associated with primary root canal treatment outcomes. The analysis particularly supported fills without radiographic voids and extending to within approximately 2 mm of the radiographic apex, while avoiding extrusion. These findings are clinically useful as process indicators but should not be treated as proof that radiographic length alone causes healing, because it may also reflect overall operator performance and anatomical difficulty [2]. A separate meta-analysis indicated that both the technical quality of root fillings and coronal restoration are relevant to treatment success [3].

Imaging and outcome definitions contribute further variability. Conventional periapical radiography permits repeated assessment at modest radiation exposure, whereas limited-field cone-beam computed tomography can reveal lesions or anatomical complications not evident on two-dimensional views. The ESE position statement recommends CBCT according to clinical indication rather than routine use in every patient [7]. The distinction between asymptomatic status, radiographic improvement, complete healing, and tooth survival is also essential. An apparently favorable short-term symptom response does not by itself establish radiographic resolution; conversely, a lesion that is decreasing in size may justify continued observation rather than premature categorization as persistent disease [1,8].

Materials and Methods

This manuscript was designed as a critical narrative evidence synthesis with an original conceptual framework for clinical quality assessment. Relevant English-language guidance and reports were identified through targeted searches of publicly accessible PubMed records, journal pages, and professional-society resources. Sources were prioritized if they addressed at least one of the following domains: treatment of apical periodontitis, irrigation and adjunctive disinfection, root filling techniques and materials, radiographic criteria of obturation quality, coronal restoration, imaging, or longitudinal healing. The core evidence set comprised the 2023 ESE S3 guideline, reviews commissioned in the same guideline program, established outcome meta-analyses, and additional clinical-methodological publications [1–10].

Journal of Advances in Drug Research

ISSN: 2230-7761
www.jadr.co.in

Volume 17, Issue 05 | Received 2.11.2026 | Accepted 7.11.2026 | Online 12.11.2026

For analytical consistency, the evidence was organized into four interrelated domains: baseline disease and tooth factors; irrigation and disinfection; obturation and coronal sealing; and patient-centered or radiographic follow-up. Strength of evidence and interpretation were separated from treatment plausibility: laboratory antimicrobial reductions were not treated as equivalent to clinical healing, and associations in nonrandomized cohorts were not interpreted as direct causal effects. Given the non-systematic search and heterogeneity of source outcomes, no pooled effect size, formal risk-of-bias score, or meta-analytic inference was calculated. There were no study participants, clinical interventions, institutional ethics approvals, or original patient observations in this manuscript.

A prospective observational protocol was then drafted as an applied research output, not a completed patient study. The proposed unit of analysis is a treated permanent tooth with diagnosed chronic apical periodontitis; multiple teeth within the same patient would require clustering adjustment. At baseline, tooth type, number of roots, lesion dimensions, previous treatment, systemic risk factors, operator, and radiographic features should be recorded. During treatment, irrigation sequence, approximate concentration, method of delivery, any activation, working length, obturation technique and material, post-obturation filling length, homogeneity and extrusion should be registered. Tooth restoration status and standardized clinical and radiographic assessment at approximately 6, 12, and 24 months are proposed to enable outcome classification. These time points are a suggested research schedule rather than observed follow-up of patients.

Results

The narrative synthesis identified a consistent distinction between fundamental standards of endodontic care and uncertain advantages of optional technologies. The ESE guideline strongly emphasizes meticulous isolation and asepsis, whereas comparisons of different irrigation concentrations, sequences, or adjunctive techniques are limited by comparatively low or very low certainty for key healing-related outcomes [1]. Published evidence for conventional irrigation regimens includes only a small number of outcome-focused trials; consequently, precise superiority rankings among commonly used solutions would be unjustified. Adjunctive disinfection strategies may improve selected technical or microbiological endpoints, but published clinical comparisons do not establish routinely superior radiographic healing relative to appropriately performed conventional irrigation [4,5,9].

The quantitative availability of evidence reported in the ESE guideline is summarized in Figure 1. For conventional irrigation comparisons, the guideline identified two randomized trials involving 212 patients for postoperative pain and one trial involving 86 patients for radiographic healing at approximately one year. For additional intracanal procedures, seven randomized trials including 636 patients addressed postoperative pain, and six randomized trials including 726 patients assessed radiographic healing; a separate cohort also contributed healing data. These figures describe sample sizes in the underlying literature, not effect sizes or evidence of equivalence or superiority [1].

Journal of Advances in Drug Research
ISSN: 2230-7761
www.jadr.co.in
Volume 17, Issue 05 | Received 2.11.2026 | Accepted 7.11.2026 | Online 12.11.2026




Figure 1. Sample sizes of comparative clinical evidence summarized by the ESE S3 guideline [1]. Study populations and endpoints differ; bar heights must not be interpreted as treatment efficacy.

Across obturation studies, currently evaluated systems show no reliable hierarchy based on long-term apical healing alone. Comparative trials have assessed gutta-percha fillings performed with several compaction techniques and sealer categories, including calcium-silicate-based materials, but the certainty and consistency of clinical outcomes do not permit the claim that a specific technique guarantees superior periapical repair. Technical assessment is more reproducible when length, voids, filling density, extrusion and restoration quality are reported independently, rather than merged into a single undocumented subjective rating [1,6].

The strongest observationally supported quality dimensions are presented in Table 1. Radiographic filling quality and the integrity of the coronal seal are complementary, not interchangeable; both can be compromised even when symptoms improve early after therapy. Preoperative lesion status and procedural complexity may confound their apparent associations with healing. Because individual patient datasets were not available, this review did not estimate odds ratios, survival probabilities, lesion-size reductions, or time-to-healing distributions for the author's practice [2,3,8].

Table 1. Integrated clinical interpretation of irrigation and obturation indicators

Domain	Measurable quality indicator	Interpretation and limitations
Asepsis / access	Dental dam use; visualization; working-length control	Foundational quality requirement; omission may raise contamination risk, but does not yield a numeric failure probability [1].
Irrigation	Solution and concentration; sequence; volume; delivery approach; complications	NaOCl–EDTA sequence may be considered. No specific regimen or activation device

Journal of Advances in Drug Research
ISSN: 2230-7761
www.jadr.co.in
Volume 17, Issue 05 Received 2.11.2026 Accepted 7.11.2026 Online 12.11.2026

Domain	Measurable quality indicator	Interpretation and limitations
		has established universal healing superiority [1,4,5].
Apical filling extent	Radiographic terminus relative to apex; unintentional extrusion	Historically associated with outcomes; interpret with anatomy and length measurement uncertainty [2].
Filling homogeneity	Presence, location and extent of radiographic voids	Absence of visible voids is associated with favorable outcomes; two-dimensional films may miss defects [2,3].
Coronal restoration	Definitive seal, marginal integrity, fractures or recurrent caries	Critical for protection against reinfection and long-term function [1,3].
Follow-up	Pain, swelling, sinus tract, function, lesion trajectory and tooth retention	Clinical improvement and radiographic healing are related but distinct endpoints [1,7,8].

Table 2 outlines a practical data dictionary for subsequent validation of a quality–outcome model. Each variable is selected to allow reproducible audit rather than to imply that its prognostic contribution has already been quantified in this article. The proposed design prioritizes separate recording of irrigation and obturation quality while accounting for baseline lesion characteristics and quality of restoration; it therefore provides a framework for more defensible multivariable analyses of later healing.

Table 2. Proposed prospective assessment framework (not observed study results)

Assessment period	Variables to document	Outcome relevance
Baseline	Age; tooth type; first treatment/retreatment; lesion dimensions; baseline imaging; symptoms; systemic conditions	Adjustment for initial risk and disease severity.
Intraoperative	Rubber dam; irrigation chemistry and method; working length; complications; obturation technique	Separate treatment exposures and adverse events.
Immediately postoperative	Root filling length and voids; extrusion; definitive restoration condition	Reproducible quality classification before follow-up.
6–24 months (proposed)	Clinical symptoms, sinus tract, functional tooth retention;	Prespecified complete healing, healing in progress, persistent

Journal of Advances in Drug Research
ISSN: 2230-7761
www.jadr.co.in
Volume 17, Issue 05 Received 2.11.2026 Accepted 7.11.2026 Online 12.11.2026

Assessment period	Variables to document	Outcome relevance
	standardized lesion assessment	disease or extraction.
Analysis plan	Blinded imaging assessment; multivariable regression; clustering by patient	Reduces measurement bias; estimates associations rather than assuming causality.




Figure 2. Proposed endodontic quality–outcome pathway. This conceptual diagram is intended for future observational validation and does not depict measured effects.

Discussion

The reviewed studies support a biologically coherent but methodologically cautious interpretation of the relationship between root canal treatment quality and clinical prognosis. Canal disinfection remains central because apical periodontitis is driven by intracanal infection; however, it is difficult to isolate the independent impact of irrigation from instrumentation, anatomy, apical enlargement and operator performance. Evidence that one activation system removes more debris or reduces bacterial counts under laboratory conditions cannot be assumed to demonstrate a corresponding increase in radiographic healing. This explains why professional recommendations distinguish basic quality principles from optional technologies that lack sufficiently consistent patient-level outcome evidence [1,4,5].

The same distinction applies to obturation. Radiographic adequacy is an informative surrogate for a carefully executed technical procedure, especially when evaluating extension and homogeneity, but cannot directly measure residual microorganisms, uninstrumented canal spaces, adhesive durability or the integrity of apical tissues. A high-quality restoration placed promptly after root canal treatment reduces the opportunity for coronal recontamination, and clinical success depends on both an internal and an external barrier. The findings of Ng et al. and Gillen et al. are therefore best interpreted as support

Journal of Advances in Drug Research
ISSN: 2230-7761
www.jadr.co.in
Volume 17, Issue 05 Received 2.11.2026 Accepted 7.11.2026 Online 12.11.2026

for coordinated technical standards rather than an isolated threshold capable of predicting individual outcomes with certainty [2,3].

Important limitations arise when comparing prior studies. Different investigations use distinct definitions of radiographic success, include primary and retreatment cases, vary in baseline lesion severity, and assess outcomes at unequal follow-up intervals. Two-dimensional imaging may underestimate lesions or fail to capture the geometry of a three-dimensional root canal filling; conversely, routinely performing CBCT in every study participant may be unjustified by radiation burden and clinical indication. A rigorous follow-up protocol should define imaging criteria in advance, use CBCT selectively when justified, and consider healing as a process that can remain incomplete at an early recall [1,7,8].

The proposed prospective framework attempts to improve methodological clarity by registering clinical and technical variables as distinct exposures. For example, an irrigation activation device should not be categorized as a "high-quality" treatment simply because it is technically sophisticated; evidence of delivered disinfection and clinical follow-up remain necessary. Likewise, radiographic root filling length should be evaluated alongside lesion status, restoration quality and the complexity of canal anatomy. To avoid biased associations, the study would need predefined inclusion criteria, calibrated image readers masked to intraoperative exposures, transparent follow-up loss accounting, and multivariable models that separate patient-level factors from tooth-level treatment features.

The present article has limitations. It is a targeted narrative synthesis rather than a registered systematic review; therefore, publication selection and qualitative interpretation are susceptible to bias. The numbers visualized in Figure 1 reflect published trial enrollment summarized by a guideline and are not pooled effect estimates. No individual patient records, microbiological measurements, standardized imaging datasets or radiographic healing outcomes were supplied for the named author. Consequently, the article cannot establish the comparative effectiveness of any locally implemented irrigation or obturation strategy. Its contribution is instead a structured translation of existing evidence into an explicit, testable clinical assessment framework.

Conclusion

In chronic apical periodontitis, effective endodontic treatment depends on the coordinated performance of aseptic isolation, chemomechanical cleaning, appropriate irrigation, technically adequate root filling, reliable coronal restoration and longitudinal monitoring. Contemporary guidance recognizes the value of sodium hypochlorite-based irrigation with EDTA while emphasizing that clinical evidence does not justify universal claims of superiority for supplementary activation procedures. The comparative literature likewise does not establish one obturation technique or sealer as consistently superior for periapical healing; filling homogeneity, appropriate extension, absence of major complications, and a durable coronal seal remain clinically relevant quality indicators.

The integrated set of indicators proposed in this manuscript offers a basis for prospective research comparing irrigation and obturation characteristics with symptomatic and

Journal of Advances in Drug Research
ISSN: 2230-7761
www.jadr.co.in
Volume 17, Issue 05 Received 2.11.2026 Accepted 7.11.2026 Online 12.11.2026

radiographic outcomes. Before any claim of original clinical effectiveness is made, this framework must be evaluated using ethically approved patient-level observations, reproducible imaging, documented follow-up and appropriate statistical adjustment. The current findings should be understood as an evidence-based synthesis and methodological proposal, not as results from a completed clinical trial.

Declarations

Ethics approval: Not applicable to this narrative review; no new research involving human participants or animals was conducted. Funding: Not reported. Competing interests: To be declared by the author before submission. Data availability: No new patient-level dataset was generated. Author contributions: To be confirmed by the author; authorship and approval of the final submitted manuscript remain the author's responsibility.

References

1. Duncan HF, Kirkevang LL, Peters OA, et al. Treatment of pulpal and apical disease: The European Society of Endodontology (ESE) S3-level clinical practice guideline. International Endodontic Journal. 2023;56(Suppl 3):238–295. https://doi.org/10.1111/iej.13974
2. Ng Y-L, Mann V, Rahbaran S, Lewsey J, Gulabivala K. Outcome of primary root canal treatment: systematic review of the literature—Part 2. Influence of clinical factors. International Endodontic Journal. 2008;41(1):6–31. https://doi.org/10.1111/j.1365-2591.2007.01323.x
3. Gillen BM, Looney SW, Gu L-S, et al. Impact of the quality of coronal restoration versus the quality of root canal fillings on success of root canal treatment: a systematic review and meta-analysis. Journal of Endodontics. 2011;37(7):895–902. https://doi.org/10.1016/j.joen.2011.04.002
4. Rossi-Fedele G, Rödig T. Effectiveness of root canal irrigation and dressing for the treatment of apical periodontitis: a systematic review and meta-analysis of clinical trials. International Endodontic Journal. 2023;56(Suppl 3):422–435. https://doi.org/10.1111/iej.13777
5. Meire MA, Bronzato JD, Bomfim RA, Gomes BPFA. Effectiveness of adjunct therapy for the treatment of apical periodontitis: a systematic review and meta-analysis. International Endodontic Journal. 2023;56(Suppl 3):455–474. https://doi.org/10.1111/iej.13838
6. Pirani C, Camilleri J. Effectiveness of root canal filling materials and techniques for treatment of apical periodontitis: a systematic review. International Endodontic Journal. 2023;56(Suppl 3):436–454. https://doi.org/10.1111/iej.13787
7. Patel S, Brown J, Semper M, Abella F, Mannocci F. European Society of Endodontology position statement: Use of cone beam computed tomography in Endodontics. International Endodontic Journal. 2019;52(12):1675–1678. https://doi.org/10.1111/iej.13187

Journal of Advances in Drug Research

ISSN: 2230-7761
www.jadr.co.in

Volume 17, Issue 05 | Received 2.11.2026 | Accepted 7.11.2026 | Online 12.11.2026

8. Rossi-Fedele G, Ng Y-L. Effectiveness of root canal treatment for vital pulps compared with necrotic pulps in the presence or absence of signs of periradicular pathosis: a systematic review and meta-analysis. International Endodontic Journal. 2023;56(Suppl 3):370–394. https://doi.org/10.1111/iej.13833
9. Immich F, Rödig T, Kanzow P, Piva E, Rossi-Fedele G. Effectiveness of Root Canal Irrigation and Dressing for the Treatment of Asymptomatic Apical Periodontitis: A Systematic Review Assessing Complementary Timeframes. Australian Endodontic Journal. 2025;51(1):209–217. https://doi.org/10.1111/aej.12918
10. Ng Y-L, Mann V, Gulabivala K. A prospective study of the factors affecting outcomes of non-surgical root canal treatment: part 2: tooth survival. International Endodontic Journal. 2011;44(7):610–625. https://doi.org/10.1111/j.1365-2591.2011.01873.x